

University of Wisconsin - Madison

MADPH-98-1063  
HAWAII-511-905-98  
VAND-TH-98-08  
AMES-HET-98-09  
June 1998

## BI-MAXIMAL MIXING OF THREE NEUTRINOS

V. Barger<sup>1</sup>, S. Pakvasa<sup>2</sup>, T.J. Weiler<sup>3</sup>, and K. Whisnant<sup>4</sup>

<sup>1</sup>*Department of Physics, University of Wisconsin, Madison, WI 53706, USA*

<sup>2</sup>*Department of Physics and Astronomy, University of Hawaii, Manoa, HI 96822, USA*

<sup>3</sup>*Department of Physics and Astronomy, Vanderbilt University, Nashville, TN 37235, USA*

<sup>4</sup>*Department of Physics and Astronomy, Iowa State University, Ames, IA 50011, USA*

### Abstract

We show that if the solar and atmospheric data are both described by maximal vacuum oscillations at the relevant mass scales then there exists a unique mixing matrix for three neutrino flavors. The solution necessarily conserves  $CP$  and automatically implies that there is no disappearance of atmospheric  $\nu_e$ , consistent with indications from the Super-Kamiokande experiment. We also investigate the consequences for three-neutrino mixing if the solar and atmospheric oscillations exhibit mixing that is large but not maximal. For non-maximal mixing  $\nu_e \leftrightarrow \nu_\tau$  and  $\nu_e \leftrightarrow \nu_\mu$  oscillations are predicted that may be observable in future long-baseline experiments.

*Introduction.* It is now understood that neutrino oscillations can describe the data on the solar neutrino deficit [1, 2, 3], the atmospheric neutrino anomaly [4, 5, 6], and the results from the LSND experiment [7], only if a sterile neutrino is introduced [8, 9, 10, 11, 12]. However, because confirmation of the LSND results awaits future experiments [13], a conservative approach is to assume that oscillations need only account for the solar and atmospheric data; also, it is possible to introduce new lepton-flavor changing operators with coefficients small enough to evade present exclusion limits, but large enough to explain the small LSND amplitude [14]. Since a comparison of the theoretical expectations [15] with the Super-Kamiokande (SuperK) atmospheric data suggests maximal  $\nu_\mu$  oscillations with mass-squared difference  $\delta m_{atm}^2 \approx 5 \times 10^{-3} \text{ eV}^2$  [5], and in light of the recent data from Super-Kamiokande [3] favoring vacuum long-wavelength oscillations [16, 17, 18] with  $\delta m_{sun}^2 \approx 10^{-10} \text{ eV}^2$  and large mixing as a description of the solar neutrino deficit, it is instructive to determine how maximal or near-maximal solar and atmospheric vacuum oscillations may be accommodated within a three-neutrino universe.

In this letter we use unitarity constraints to derive conditions on the three-neutrino mixing matrix under the assumption that both solar and atmospheric neutrinos undergo maximal mixing *in vacuum*. By maximal mixing we mean that the disappearance probabilities are equivalent to those for maximal two-neutrino mixing at the relevant mass scales. We find that there exists a unique three-neutrino mixing solution, up to trivial sign ambiguities, which conserves  $CP$  and has no oscillations of atmospheric  $\nu_e$ , consistent with indications from the Super-Kamiokande experiment [5]. This mixing matrix predicts that solar  $\nu_e$  oscillate maximally into equal numbers of  $\nu_\mu$  and  $\nu_\tau$ . We also investigate the consequences for three-neutrino mixing if the solar and atmospheric oscillations are not maximal. We find that for near-maximal solar and atmospheric neutrino mixing the  $\nu_\mu \rightarrow \nu_e$  and  $\nu_e \rightarrow \nu_\tau$  oscillation amplitudes for atmospheric and long-baseline experiments are approximately equal, and are limited by the deviation of the solar  $\nu_e$  disappearance amplitude from unity. Also, for any non-maximal solar and atmospheric neutrino mixing there must exist  $\nu_e \rightarrow \nu_\tau$  oscillations that may be visible in future long-baseline experiments.

*Oscillation probabilities.* We begin our analysis with the survival probability for neutrino oscillations in a vacuum [19]

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - 4 \sum_{k < j} P_{\alpha j} P_{\alpha k} \sin^2 \Delta_{jk}, \quad (1)$$

where  $P_{\alpha j} \equiv |U_{\alpha j}|^2$  is the probability to find the  $\alpha$ -flavor state in the  $j$ th mass state (or vice versa),  $U$  is the neutrino mixing matrix (in the basis where the charged-lepton mass matrix is diagonal),  $\Delta_{jk} \equiv \delta m_{jk}^2 L / 4E = 1.27(\delta m_{jk}^2 / \text{eV}^2)(L/\text{km})/(E/\text{GeV})$ ,  $\delta m_{jk}^2 \equiv m_j^2 - m_k^2$ , and the sum is over all  $j$  and  $k$ , subject to  $k < j$ . The matrix elements  $U_{\alpha j}$  are the mixings between the flavor ( $\alpha = e, \mu, \tau$ ) and the mass ( $j = 1, 2, 3$ ) eigenstates. We assume a neutrino mass spectrum with mass eigenvalues  $m_1, m_2 \ll m_3$ , where the solar oscillations are driven by  $\Delta_{21} \equiv \Delta_{sun}$  and the atmospheric oscillations are driven by  $\Delta_{31} \simeq \Delta_{32} \equiv \Delta_{atm}$ . However, only the form of the neutrino mass matrix depends on the mass hierarchy assumption. All of our other conclusions require only the more general assumption that  $|\Delta_{sun}| = |\Delta_{21}| \ll |\Delta_{31}| \simeq |\Delta_{32}| = |\Delta_{atm}|$ , which also could apply, e.g., to the case  $m_3 < m_1 \simeq m_2$ , i.e., the solar neutrino vacuum oscillations are driven by a small mass splitting between the two larger masses.

The off-diagonal oscillation probabilities of this model are

$$P(\nu_e \rightarrow \nu_\mu) = 4 P_{e3} P_{\mu3} \sin^2 \Delta_{atm} - 4 \text{Re}\{U_{e1} U_{e2}^* U_{\mu1}^* U_{\mu2}\} \sin^2 \Delta_{sun} - 2 J \sin 2\Delta_{sun}, \quad (2)$$

$$P(\nu_e \rightarrow \nu_\tau) = 4 P_{e3} P_{\tau3} \sin^2 \Delta_{atm} - 4 \text{Re}\{U_{e1} U_{e2}^* U_{\tau1}^* U_{\tau2}\} \sin^2 \Delta_{sun} + 2 J \sin 2\Delta_{sun}, \quad (3)$$

$$P(\nu_\mu \rightarrow \nu_\tau) = 4 P_{\mu3} P_{\tau3} \sin^2 \Delta_{atm} - 4 \text{Re}\{U_{\mu1} U_{\mu2}^* U_{\tau1}^* U_{\tau2}\} \sin^2 \Delta_{sun} - 2 J \sin 2\Delta_{sun}, \quad (4)$$

where the  $CP$ -violating “Jarlskog invariant” [20] is  $J = \text{Im}\{U_{e2} U_{e3}^* U_{\mu2}^* U_{\mu3}\}$ . The  $CP$ -odd term changes sign under reversal of the oscillating flavors. The solar amplitudes, quartic in the  $U_{\alpha j}$ ’s, may themselves be expressed in terms of the  $P_{\alpha j}$  [21, 22], as may the  $CP$ -odd amplitude  $J$  [22, 23].

We note that the  $CP$ -violating probability at the atmospheric scale is suppressed to order  $\delta m_{sun}^2 / \delta m_{atm}^2$ , the leading term having canceled in the sum over the two light-mass states. Thus,  $P(\nu_\alpha \rightarrow \nu_\beta) = P(\nu_\beta \rightarrow \nu_\alpha)$  at the atmospheric scale. With MSW-enhanced solutions to the solar deficit,  $\sin 2\Delta_{sun}$  is effectively replaced by its average value of zero since  $\Delta_{sun} \gg 1$  in this case. Only for the vacuum long-wavelength solution to the solar neutrino anomaly is there hope to measure the  $CP$ -odd term.

We will use “amplitude” to denote the coefficients of the oscillating factors. The amplitudes of interest are:

$$A_{sun}^{e\ell} = 4 P_{e1} P_{e2} \leq (1 - P_{e3})^2, \quad (5)$$

$$A_{atm}^{\mu\mu} = 4 P_{\mu3} (P_{\mu1} + P_{\mu2}) = 4 P_{\mu3} (1 - P_{\mu3}), \quad (6)$$

$$A_{atm}^{e\ell} = 4 P_{e3} (1 - P_{e3}), \quad (7)$$

$$A_{atm}^{\mu e} = 4 P_{e3} P_{\mu3} \leq (1 - P_{\tau3})^2, \quad (8)$$

$$A_{atm}^{\mu\tau} = 4 P_{\mu3} P_{\tau3} \leq (1 - P_{e3})^2, \quad (9)$$

$$A_{atm}^{e\tau} = 4 P_{e3} P_{\tau3} \leq (1 - P_{\mu3})^2. \quad (10)$$

In a three-neutrino model

$$A_{atm}^{\mu\mu} = A_{atm}^{\mu e} + A_{atm}^{\mu\tau}, \quad \text{and} \quad (11)$$

$$A_{atm}^{e\ell} = A_{atm}^{\mu e} + A_{atm}^{e\tau}. \quad (12)$$

The equality in Eq. (5) results from the unitarity condition  $U_{\mu1}^* U_{\mu2} + U_{\tau1}^* U_{\tau2} = -U_{e1}^* U_{e2}$ . The inequalities in Eqs. (5), (8)–(10) are obtained by maximizing the amplitude under the unitarity constraints

$$P_{e3} + P_{\mu3} + P_{\tau3} = 1, \quad \text{and} \quad (13)$$

$$P_{e1} + P_{e2} + P_{e3} = 1. \quad (14)$$

Thus, the amplitudes depend on just three independent  $P_{\alpha j}$ ’s, and the atmospheric amplitudes depend on just two of the probabilities chosen from  $P_{e3}$ ,  $P_{\mu3}$ , and  $P_{\tau3}$ .

*Maximal solar and atmospheric oscillations.* We first explore a solution for which both solar  $\nu_e$  and atmospheric  $\nu_\mu$  data can be described by maximal oscillations at the relevant  $\delta m^2$  scales, as suggested by the SuperK data [3, 5]. Maximal atmospheric oscillations require

$P_{\mu 3} = \frac{1}{2}$ . If solar oscillations are also required to be maximal, then  $P_{e3} = 0$  and  $P_{e1} = P_{e2} = \frac{1}{2}$ . Up to trivial sign changes, unitarity then uniquely specifies the remaining elements of  $U$ :

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix} = \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} & 0 \\ \frac{1}{2} & \frac{1}{2} & -\frac{1}{\sqrt{2}} \\ \frac{1}{2} & \frac{1}{2} & \frac{1}{\sqrt{2}} \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}. \quad (15)$$

With the mixing matrix of Eq. (15), the off-diagonal oscillation probabilities in Eqs. (2)–(4) are given by

$$P(\nu_e \rightarrow \nu_\mu) = \frac{1}{2} \sin^2 \Delta_{sun}, \quad (16)$$

$$P(\nu_e \rightarrow \nu_\tau) = \frac{1}{2} \sin^2 \Delta_{sun}, \quad (17)$$

$$P(\nu_\mu \rightarrow \nu_\tau) = \sin^2 \Delta_{atm} - \frac{1}{4} \sin^2 \Delta_{sun}. \quad (18)$$

The zero  $U_{e3}$  element implies that  $CP$  is conserved,  $J$  is zero, and the elements of  $U$  are real. It also implies that the only oscillations at the  $\Delta_{atm}$  scale are  $\nu_\mu \rightarrow \nu_\tau$ . The solar  $\nu_e$  oscillations are 50% into  $\nu_\mu$  and 50% into  $\nu_\tau$ , although the oscillation flavors are not distinguishable because they have the same neutral current interactions and there is not enough energy to produce charge current events; however, detection of neutrinos from active galactic nuclei and gamma ray burst sources could distinguish the  $\tau$ -neutrino flavor from  $\nu_e$  and  $\nu_\mu$  [24]. The mixing matrix for this bi-maximal scenario is similar to that obtained in Refs. [25] and [26], which give maximal solar oscillations ( $A_{sun}^{e\ell} = 1$ ) and large but not maximal mixing of atmospheric neutrinos ( $A_{atm}^{\mu\mu} = \frac{8}{9}$ ). All of these scenarios also have  $U_{e3} = 0$ , which implies maximal solar mixing and only  $\nu_\mu \rightarrow \nu_\tau$  oscillations at the  $\Delta_{atm}$  scale.

A Majorana mass matrix or a hermitian Dirac mass matrix can be diagonalized by a single unitary matrix. For these two cases, we may therefore invert the process to obtain  $M$  in the flavor basis from the mass eigenvalues  $m_j$  and the mixing matrix  $U$ . For the Majorana case  $M_{Maj} = U M_{diag} U^T$ , while for the hermitian Dirac case  $M_{Dirac} = U M_{diag} U^\dagger$ . The individual matrix elements are  $M_{\alpha\beta} = \sum_j m_j U_{\alpha j} U_{\beta j}$  and  $M_{\alpha\beta} = \sum_j m_j U_{\alpha j} U_{\beta j}^*$ , respectively. For real  $U$ , as here, there is no difference between these two cases. Reconstruction of the mass matrix from  $U$  in Eq. (15) gives

$$M = m \left[ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & -1 \\ 0 & -1 & 1 \end{pmatrix} + \begin{pmatrix} 2\epsilon & \delta & \delta \\ \delta & \epsilon & \epsilon \\ \delta & \epsilon & \epsilon \end{pmatrix} \right], \quad (19)$$

in the  $(\nu_e, \nu_\mu, \nu_\tau)$  basis, where the parameters of  $M$  are related to the mass eigenvalues as

$$m = m_3/2, \quad (20)$$

$$\epsilon = (m_2 + m_1)/4m, \quad (21)$$

$$\delta = \sqrt{2}(m_1 - m_2)/4m. \quad (22)$$

Individual field redefinitions  $\psi_\alpha \rightarrow -\psi_\alpha$  and  $\psi_j \rightarrow -\psi_j$  allow one to change the signs of individual rows and columns of  $M$ . We remind the reader that the form of  $M$  derived here applies in the basis where the charged-lepton mass matrix is diagonal.

The two relevant mass-squared differences are

$$\delta m_{atm}^2 \simeq 4 m^2, \quad \delta m_{sun}^2 = 8 \sqrt{2} \delta \epsilon m^2. \quad (23)$$

The atmospheric oscillation scale requires  $m \approx 0.035$  eV. The vacuum long-wavelength solar solution then requires  $|\delta \epsilon| \approx 10^{-8}$ . Both  $\delta$  and  $\epsilon$  must be nonzero to generate a nonzero  $\delta m_{sun}^2$ . We reiterate that only the form of the mass matrix in Eq. (19) depends on the assumption  $m_1, m_2 \ll m_3$ ; the oscillation probabilities hold for the more general relation  $|\Delta_{sun}| \ll |\Delta_{atm}|$ .

*Near-maximal mixing.* We next consider having both solar and atmospheric neutrino mixing close to maximal ( $A_{atm}^{\mu\mu} \approx 1$  and  $A_{sun}^{e\ell} \approx 1$ ). Parametrizing the real mixing matrix in terms of Euler angles

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix} = \begin{pmatrix} c_1 c_3 & -c_1 s_3 & s_1 \\ s_1 c_2 c_3 + s_2 s_3 & -s_1 c_2 s_3 + s_2 c_3 & -c_1 c_2 \\ -s_1 s_2 c_3 + c_2 s_3 & s_1 s_2 s_3 + c_2 c_3 & c_1 s_2 \end{pmatrix} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}, \quad (24)$$

where  $c_j \equiv \cos \theta_j$  and  $s_j \equiv \sin \theta_j$ , the maximal solar and atmospheric mixing of Eq. (15) corresponds to  $\theta_1 = 0$ ,  $\theta_2 = \frac{\pi}{4}$ , and  $\theta_3 = \frac{\pi}{4}$ . Then for near-maximal mixing we take

$$\theta_1 = \epsilon_1, \quad \theta_2 = \frac{\pi}{4} + \epsilon_2, \quad \theta_3 = \frac{\pi}{4} + \epsilon_3, \quad (25)$$

where the  $\epsilon_j$  are small. The corresponding oscillation amplitudes are

$$A_{sun}^{e\ell} \simeq 1 - 2\epsilon_1^2 - 4\epsilon_3^2, \quad (26)$$

$$A_{atm}^{\mu\mu} \simeq 1 - 4\epsilon_2^2, \quad (27)$$

$$A_{atm}^{\mu e} \simeq A_{atm}^{e\tau} \simeq \frac{1}{2} A_{atm}^{e\ell} \simeq 2\epsilon_1^2. \quad (28)$$

Here atmospheric  $\nu_\mu$  disappearance is determined solely by  $\epsilon_2$ . The solar  $\nu_e$  disappearance and atmospheric and long-baseline  $\nu_\mu \rightarrow \nu_e$  and  $\nu_e \rightarrow \nu_\tau$  oscillation amplitudes must satisfy the conditions

$$\frac{1}{2} A_{atm}^{e\ell} \simeq A_{atm}^{\mu e} \simeq A_{atm}^{e\tau} \lesssim 1 - A_{sun}^{e\ell}. \quad (29)$$

Hence for near-maximal mixing,  $\nu_\mu \rightarrow \nu_e$  and  $\nu_e \rightarrow \nu_\tau$  oscillations in atmospheric and long-baseline experiments are small and approximately equal, and they are constrained by how close the solar mixing is to maximal. The atmospheric  $\nu_\mu$  disappearance amplitude is independent of the other oscillation amplitudes in this limit.

*Non-maximal mixing.* We also investigate the more general case of large but non-maximal solar neutrino mixing, which corresponds to nonzero  $U_{e3}$ , and large but non-maximal atmospheric neutrino mixing. A geometric expression for  $J$  [20, 22] equates its magnitude to the area of a triangle with sides which may be taken to be  $|U_{e3}U_{\mu 3}|$ ,  $|U_{e2}U_{\mu 2}|$ , and  $|U_{e1}U_{\mu 1}|$ . Since  $|U_{e3}|$  is known to be small, we use Eq.(15) as an approximate guide for  $U$  and obtain  $|J| \lesssim |U_{e3}|/8$  as the order of magnitude for the area of the triangle. This bound on  $|J|$  is sufficiently small that we will set  $|J|$  to zero and continue to work with real  $U$  and three independent parameters in the mixing matrix. If the oscillation amplitudes  $A_{sun}^{e\ell}$  and  $A_{atm}^{\mu\mu}$

are determined, e.g., from experiment, then there remains only one free parameter, e.g.,  $A_{atm}^{e\ell}$ . Although pure  $\nu_\mu \leftrightarrow \nu_e$  oscillations of atmospheric neutrinos are strongly disfavored, modest amounts of  $\nu_e$  disappearance have not yet been ruled out. Therefore it is worthwhile to consider the effects of a nonzero  $U_{e3}$  when the approximations of the previous section are no longer valid.

Given the inputs  $A_{sun}^{e\ell}$ ,  $A_{atm}^{\mu\ell}$ , and  $A_{atm}^{e\ell}$ , the three independent parameters of the matrix  $U$  can be determined up to some two-fold ambiguities, only one of which has phenomenological consequences for vacuum oscillations. For example, knowing the atmospheric  $\nu_\mu$  disappearance amplitude in Eq. (6) determines  $U_{\mu 3} = \pm\sqrt{P_{\mu 3}}$ :

$$P_{\mu 3} = \frac{1}{2} \left[ 1 \mp \sqrt{1 - A_{atm}^{\mu\ell}} \right], \quad (30)$$

where the  $- (+)$  means that  $\nu_\mu$  is more closely associated with  $\nu_2 (\nu_3)$ . Knowing the atmospheric  $\nu_e$  disappearance amplitude in Eq. (7) determines  $U_{e3} = \pm\sqrt{P_{e3}}$ :

$$P_{e3} = \frac{1}{2} \left[ 1 - \sqrt{1 - A_{atm}^{e\ell}} \right]. \quad (31)$$

In Eq. (31) we have chosen the solution which gives  $P_{e3} \leq \frac{1}{2}$  since  $P_{e3} > \frac{1}{2}$  implies that the solar  $\nu_e$  disappearance amplitude  $A_{sun}^{e\ell}$  must be less than  $\frac{1}{4}$ , which for the long-wavelength solar solution is disfavored by the data. Then the unitarity relation in Eq. (13) yields

$$P_{\tau 3} = \frac{1}{2} \left[ \sqrt{1 - A_{atm}^{e\ell}} \pm \sqrt{1 - A_{atm}^{\mu\ell}} \right], \quad (32)$$

where the  $\pm$  in Eq. (32) is correlated with the  $\mp$  in Eq. (30). Also, Eq. (5) and the unitarity relation in Eq. (14) imply

$$P_{e1} = \frac{1}{2} \left[ 1 - P_{e3} + \sqrt{(1 - P_{e3})^2 - A_{sun}^{e\ell}} \right], \quad (33)$$

$$P_{e2} = \frac{1}{2} \left[ 1 - P_{e3} - \sqrt{(1 - P_{e3})^2 - A_{sun}^{e\ell}} \right], \quad (34)$$

where we have assumed that  $\nu_e$  is more closely associated with  $\nu_1$  than with  $\nu_2$  (the other possibility, that  $\nu_e$  is more closely associated with  $\nu_2$ , has similar vacuum oscillation phenomenology). In principle, there is a constant term in the solar oscillation probability  $2 P_{e3}(1 - P_{e3}) = \frac{1}{2} A_{atm}^{e\ell}$ , from averaging of the  $\sin^2 \Delta_{atm}$  term. However, if  $A_{atm}^{e\ell}$  is small, as suggested by the non-observation of  $\nu_e$  oscillations in the atmospheric data, this term can be ignored to first approximation [27]. The remaining elements of  $U$  may also be determined from unitarity (e.g., by using the values of  $P_{\mu 3}$ ,  $P_{e3}$ ,  $P_{\tau 3}$ ,  $P_{e1}$ , and  $P_{e2}$  and the form of Eq. (24) to determine the three Euler mixing angles).

Once the oscillation parameters are fixed (up to the two-fold ambiguity in Eq. (30)), other oscillation amplitudes may be calculated; for example, Eqs. (30) and (31) combined with Eq. (8) yield

$$A_{atm}^{\mu e} = \left[ 1 - \sqrt{1 - A_{atm}^{e\ell}} \right] \left[ 1 \mp \sqrt{1 - A_{atm}^{\mu\ell}} \right], \quad (35)$$

where the  $\mp$  is correlated with the  $\mp$  in Eq. (30). Contours of  $A_{atm}^{\mu e}$  in the  $(A_{atm}^{\mu\mu}, A_{atm}^{e\ell})$  plane are shown in Figs. 1(a) and 1(b). Once  $A_{atm}^{\mu\mu}$  and  $A_{atm}^{e\ell}$  are measured, the value of  $A_{atm}^{\mu e}$  is predicted.

In principle the three inputs can be determined by the solar and atmospheric neutrino data. For instance,  $A_{sun}^{e\ell}$  is directly determined from vacuum long-wavelength fits to the solar neutrino spectrum. Also, the two independent atmospheric oscillation amplitudes can be found from measurements of

$$N_\mu/N_\mu^o = (1 - \langle S \rangle A_{atm}^{\mu\mu}) + r \langle S \rangle A_{atm}^{\mu e}, \quad (36)$$

and

$$N_e/N_e^o = (1 - \langle S \rangle A_{atm}^{e\ell}) + r^{-1} \langle S \rangle A_{atm}^{\mu e}, \quad (37)$$

where  $N_e^o$  and  $N_\mu^o$  are the expected number of atmospheric  $e$  and  $\mu$  events,  $r \equiv N_e^o/N_\mu^o \sim \frac{1}{2}$ , and  $\langle S \rangle$  is  $\sin^2 \Delta_{atm}$  averaged over the appropriate atmospheric neutrino spectrum.

The existence of a real solution to Eqs. (33) and (34) requires

$$P_{e3} \leq 1 - \sqrt{A_{sun}^{e\ell}}. \quad (38)$$

The limit in Eq. (38) becomes stringent when  $A_{sun}^{e\ell}$  is close to unity. Current data suggest  $A_{sun}^{e\ell} \geq 0.6$ , which gives  $P_{e3} \leq 0.23$  and hence  $A_{atm}^{e\ell} \leq 0.7$ . These limits will improve as more solar and atmospheric data become available. As a check, note that in Eq. (38) maximal mixing of solar neutrinos ( $A_{sun}^{e\ell} = 1$ ) automatically leads to  $|U_{e3}| = A_{atm}^{\mu e} = A_{atm}^{e\tau} = 0$ , i.e., all atmospheric oscillations are  $\nu_\mu \rightarrow \nu_\tau$ . From our approximate bound for  $|J|$  when  $U_{e3}$  is small, we obtain  $|J| \lesssim 0.06$ , which verifies that  $CP$ -violation effects will be small.

Another limit on  $P_{e3}$  comes from the CHOOZ reactor experiment [28] that measures  $\bar{\nu}_e$  disappearance

$$A_{atm}^{e\ell} \lesssim 0.2, \quad (39)$$

which applies for  $\delta m_{atm}^2 \gtrsim 2 \times 10^{-3} \text{ eV}^2$ . When combined with Eq. (31), Eq. (39) gives the bound

$$P_{e3} \lesssim 0.05. \quad (40)$$

However, for  $\delta m_{atm}^2 < 10^{-3} \text{ eV}^2$  there is no limit at all. The proposed experiment to detect reactor anti-neutrinos in the BOREXINO detector [29] can test a value of  $A_{atm}^{e\ell}$  as low as 0.4 (corresponding to a limit  $P_{e3} \leq 0.18$ ) for  $\delta m_{atm}^2 \gtrsim 10^{-6} \text{ eV}^2$ .

The new physics predicted by the non-maximal mixing model is  $\nu_e \rightarrow \nu_\tau$  oscillations with leading probability (i.e., ignoring oscillations involving  $\Delta_{sun}$ )

$$P(\nu_e \rightarrow \nu_\tau) = 4 P_{e3} P_{\tau 3} \sin^2 \Delta_{atm}, \quad (41)$$

which from Eqs. (31), (32), and (10) has an oscillation amplitude

$$A_{atm}^{e\tau} = \left[ 1 - \sqrt{1 - A_{atm}^{e\ell}} \right] \left[ \sqrt{1 - A_{atm}^{e\ell}} \pm \sqrt{1 - A_{atm}^{\mu\mu}} \right], \quad (42)$$

where the  $\pm$  correlates with the  $\mp$  in Eq. (31). As with  $A_{atm}^{\mu e}$ ,  $A_{atm}^{e\tau}$  is completely determined once  $A_{atm}^{\mu\mu}$  and  $A_{atm}^{e\ell}$  are known, modulo the two-fold ambiguity in Eq. (30). Contours of  $A_{atm}^{e\tau}$

in the  $(A_{atm}^{\mu\mu}, A_{atm}^{e\ell})$  plane are shown in Figs. 1(c) and 1(d). We note that the approximate equalities involving the atmospheric oscillation amplitudes in Eq. (29) can also be derived by taking the  $A_{atm}^{\mu\mu} \rightarrow 1$ ,  $A_{atm}^{e\ell} \rightarrow 0$  limit in Eqs. (35) and (42).

The  $\nu_e \rightarrow \nu_\tau$  oscillations could be observed by long-baseline neutrino experiments with proposed high intensity muon sources [30, 10, 12], which can also make precise measurements of  $\nu_\mu \rightarrow \nu_e$ ,  $\nu_\mu \rightarrow \nu_\tau$ , and  $\nu_e \rightarrow \nu_\mu$  oscillations. Sensitivity to  $A_{atm}^{e\tau}(\delta m_{atm}^2/\text{eV}^2)^2 > 2.5 \times 10^{-9}$  is expected [30] for the parameter ranges of interest here. The combination of all measurements (solar, atmospheric, and long-baseline) would serve as a consistency check on the model.

*Other consequences.* Because the highest mass scale is  $\sqrt{\delta m_{atm}^2} \approx 0.07$  eV, there is little neutrino contribution to hot dark matter in this model [31, 32]. Such a mass can be generated for Majorana neutrinos by the see-saw mechanism with a heavy Majorana scale of about  $5 \times 10^{14}$  GeV (radiative corrections may reduce this value by about a factor of two [33]) if the Dirac mass is of order the top mass. Also, if  $m_1 \ll m_2 < m_3$  in this model, then  $m_2/m_3 \sim 10^{-4}$  is approximately equal to  $(m_c/m_t)^2$ , as predicted in some see-saw models [33]. The small value of the mass matrix element  $M_{\nu_e \nu_e} \approx 10^{-5}$  eV means that the model has an unmeasurably small contribution to neutrinoless double-beta decay [34]. The small values of  $U_{e3}$  and  $m_3$  mean that there are no observable consequences for tritium beta-decay [35]. However, the heaviest mass does provide a relic neutrino target for a mechanism that may generate the cosmic ray air showers observed above  $\gtrsim 10^{20}$  eV [36].

*Discussion.* More precise measurements of the spectrum shape and day-night effects by SuperK and the Sudbury Neutrino Observatory (SNO) [37] could determine if the solar oscillations are vacuum long-wavelength. Measurements of a seasonal variation of the flux of solar  $^7\text{Be}$  and  $pep$  neutrinos [38], which can be measured in the BOREXINO experiment [29], would confirm that the solar oscillations are vacuum long-wavelength. The HELLAZ experiment [39] plans to measure the solar  $pp$  neutrino spectrum, which would also be useful in establishing the vacuum long-wavelength solution. All of these solar experiments should also be able to pin down the value  $A_{sun}^{e\ell}$  in the vacuum long-wavelength scenario.

Further measurements of atmospheric neutrinos will more precisely determine the amplitudes  $A_{atm}^{\mu\mu}$  and  $A_{atm}^{e\ell}$ , completely fixing (modulo a two-fold ambiguity) all oscillation amplitudes in atmospheric and long-baseline experiments. The MINOS experiment [40] is expected to test for  $\nu_\mu \rightarrow \nu_e$  and  $\nu_\mu \rightarrow \nu_\tau$  oscillations for  $\delta m_{atm}^2 > 10^{-3}$  eV<sup>2</sup>. Together the solar and atmospheric measurements could put strong limits on  $U_{e3}$ , which governs the amount of  $\nu_e \rightarrow \nu_\tau$  mixing that could be seen in future long-baseline experiments such as those discussed for the proposed muon collider at Fermilab [30]. For near-maximal solar and atmospheric neutrino mixing the  $\nu_\mu \rightarrow \nu_e$  and  $\nu_e \rightarrow \nu_\tau$  oscillation amplitudes for atmospheric and long-baseline experiments are approximately equal, and are limited by the deviation of the solar  $\nu_e$  disappearance amplitude from unity.

*Acknowledgements.* We thank John Learned for stimulating discussions on mixing matrices with maximal mixing and X. Tata for discussions on see-saw mass matrices. This work was supported in part by the U.S. Department of Energy, Division of High Energy Physics, under Grants No. DE-FG02-94ER40817, No. DE-FG05-85ER40226, and No. DE-FG02-95ER40896, and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation and the Vanderbilt University Research Council.



## References

- [1] J.N. Bahcall and M.H. Pinsonneault, Rev. Mod. Phys. **67**, 781 (1995); J.N. Bahcall, S. Basu, and M.H. Pinsonneault, astro-ph/9805135.
- [2] B.T. Cleveland *et al.*, Nucl. Phys. B (Proc. Suppl.) **38**, 47 (1995); Kamiokande collaboration, Y. Fukuda *et al.*, Phys. Rev. Lett, **77**, 1683 (1996); GALLEX Collaboration, W. Hampel *et al.*, Phys. Lett. **B388**, 384 (1996); SAGE collaboration, J.N. Abdurashitov *et al.*, Phys. Rev. Lett. **77**, 4708 (1996).
- [3] Super-Kamiokande Collaboration, talk by Y. Suzuki at *Neutrino-98*, Takayama, Japan, June 1998.
- [4] Kamiokande collaboration, K.S. Hirata *et al.*, Phys. Lett. **B280**, 146 (1992); Y. Fukuda *et al.*, Phys. Lett. **B335**, 237 (1994); IMB collaboration, R. Becker-Szendy *et al.*, Nucl. Phys. Proc. Suppl. **38B**, 331 (1995); Soudan-2 collaboration, W.W.M. Allison *et al.*, Phys. Lett. **B391**, 491 (1997).
- [5] Super-Kamiokande Collaboration, talk by T. Kajita at *Neutrino-98*, Takayama, Japan, June 1998; Y. Fukuda *et al.*, hep-ex/9803006 and Phys. Lett. **B**, to appear; hep-ex/9805006; see also talk by E. Kearns at the ITP Conference on Solar Neutrinos: News About SNUS, Santa Barbara, December 1997, and hep-ex/9803007.
- [6] J.G. Learned, S. Pakvasa, and T.J. Weiler, Phys. Lett. **B207**, 79 (1988); V. Barger and K. Whisnant, Phys. Lett. **B209**, 365 (1988); K. Hidaka, M. Honda, and S. Midorikawa, Phys. Rev. Lett. **61**, 1537 (1988).
- [7] Liquid Scintillator Neutrino Detector (LSND) collaboration, C. Athanassopoulos *et al.*, Phys. Rev. Lett. **75**, 2650 (1995); *ibid.* **77**, 3082 (1996); nucl-ex/9706006; talk by H. White at *Neutrino-98*, Takayama, Japan, June 1998.
- [8] D.O. Caldwell and R.N. Mohapatra, Phys. Rev. **D 48**, 3259 (1993); J.T. Peltoniemi and J.W.F. Valle, Nucl. Phys. **B 406**, 409 (1993); R. Foot and R.R. Volkas, Phys. Rev. **D 52**, 6595 (1995); J.J. Gomez-Cadenas and M.C. Gonzalez-Garcia, Z. Phys. **C 71**, 443 (1996); S.M. Bilenky, C. Giunti, and W. Grimus, hep-ph/9711416; R.N. Mohapatra, hep-ph/9711444; D.O. Caldwell, hep-ph/9804367.
- [9] S. M. Bilenky, C. Giunti and W. Grimus, Eur. Phys. J. C1, 247 (1998) and hep-ph/9607372.
- [10] V. Barger, K. Whisnant, and T.J. Weiler, hep-ph/9712495 and Phys. Lett. **B**, to appear.
- [11] S.C. Gibbons, R.N. Mohapatra, S. Nandi, and A. Raychaudhuri, hep-ph/9803299.
- [12] V. Barger, S. Pakvasa, T.J. Weiler, and K. Whisnant, hep-ph/9806328.
- [13] The LSND results will be tested by the KARMEN experiment, talk by B. Armbruster at 33rd Rencontres de Moriond: Electroweak Interactions and Unified Theories, Les Arcs, France, March 1998, and talk by B. Zeitnitz at *Neutrino-98*, Takayama, Japan, June 1998, and also by the BooNE experiment, E. Church *et al.*, nucl-ex/9706011.

- [14] L.M. Johnson and D. McKay, hep-ph/9805311.
- [15] G. Barr, T.K. Gaisser, and T. Stanev, Phys. Rev. **D 39**, 3532 (1989); M. Honda, T. Kajita, K. Kasahara, and S. Midorikawa, Phys. Rev. **D52**, 4985 (1995); V. Agrawal, T.K. Gaisser, P. Lipari, and T. Stanev, Phys. Rev. **D 53**, 1314 (1996); T.K. Gaisser *et al.*, Phys. Rev. **D 54**, 5578 (1996).
- [16] V. Barger, R.J.N. Phillips, and K. Whisnant, Phys. Rev. **D 24**, 538 (1981).
- [17] S.L. Glashow and L.M. Krauss, Phys. Lett. **B190**, 199 (1987).
- [18] A. Acker, S. Pakvasa, and J. Pantaleone, Phys. Rev. **D 43**, 1754 (1991); V. Barger, R.J.N. Phillips, and K. Whisnant, Phys. Rev. Lett. **69**, 3135 (1992); P.I. Krastev and S.T. Petcov, Phys. Lett. **B 285**, 85 (1992); Phys. Rev. Lett. **72**, 1960 (1994); Phys. Rev. **D 53**, 1665 (1996).
- [19] V. Barger, K. Whisnant, D. Cline, and R.J.N. Phillips, Phys. Lett. **B93**, 194 (1980).
- [20] C. Jarlskog, Z. Phys. **C29**, 491 (1985); Phys. Rev. **D35**, 1685 (1987).
- [21] C.W. Kim and A. Pevsner, *Neutrinos in Physics and Astrophysics*, Hardwood Academic Pub., 1993.
- [22] D.J. Wagner and T.J. Weiler, Phys. Rev. **D**, to appear, and hep-ph/9801327.
- [23] C. Hamzaoui, Phys. Rev. Lett. **61**, 35 (1988); C. Jarlskog, in *CP Violation*, ed. by C. Jarlskog (World Scientific, Singapore, 1989).
- [24] J.G. Learned and S. Pakvasa, Astropart. Phys. **3**, 267 (1995); F. Halzen and D. Saltzberg, hep-ph/9804354.
- [25] H. Fritzsch and Z. Xing, Phys. Lett. **B 372**, 265 (1996).
- [26] M. Fukugita, M. Tanimoto, and T. Yanagida, hep-ph/9709388.
- [27] The effects of nonzero  $U_{e3}$  on the solar fit are considered by P. Osland and G. Vigdel, hep-ph/9806339.
- [28] CHOOZ collaboration, M. Apollonio *et al.*, hep-ex/9711002.
- [29] BOREXINO Collaboration, C. Arpesella *et al.*, “INFN Borexino proposal,” Vols. I and II, edited by G. Bellini, R. Raghavan *et al.* (Univ. of Milan, 1992); J. Benziger, F.P. Calaprice *et al.*, “Proposal for Participation in the Borexino Solar Neutrino Experiment,” (Princeton University, 1996).
- [30] S. Geer, hep-ph/9712290.
- [31] E.W. Kolb and M.S. Turner, *The Early Universe* (Addison-Wesley, Reading, 1990).
- [32] For a recent discussion see J. Primack, astro-ph/9707285.

- [33] S.A. Bludman, D.C. Kennedy, and P.G. Langacker, Phys. Rev. **D 45**, 1810 (1992).
- [34] H.V. Klapdor-Kleingrothaus, hep-ph/9802007; J. Phys. **G24**, 483 (1998); talk at *Neutrino-98*, Takayama, Japan, June 1998.
- [35] A.I. Belesev *et al.*, Phys. Lett. **B350**, 263 (1995); V.M. Lobashev *et al.*, in *Proc. of Neutrino '96*, ed. by K. Enqvist, K. Huitu, and J. Maalampi (World Scientific, Singapore, 1997).
- [36] T.J. Weiler, hep-ph/9710431, to appear in Astropart. Phys.
- [37] SNO Collaboration, E. Norman *et al.*, in proc. of *The Fermilab Conference: DPF 92*, November 1992, Batavia, IL, ed. by C. H. Albright, P.H. Kasper, R. Raja, and J. Yoh (World Scientific, Singapore, 1993), p. 1450.
- [38] V. Barger, R.J.N. Phillips, and K. Whisnant, Phys. Rev. Lett. **65**, 3084 (1990); S. Pakvasa and J. Pantaleone, Phys. Rev. Lett. **65**, 2479 (1990); A. Acker, Ph.D. thesis, Dec. 1992, UH-511-759-92. For a recent discussion, see B. Faid, G.L. Fogli, E. Lisi, and D. Montanino, hep-ph/9805293.
- [39] HELLAZ Collaboration, F. Arzarello *et al.*, CERN-LAA/94-19.
- [40] MINOS Collaboration, “Neutrino Oscillation Physics at Fermilab: The NuMI-MINOS Project,” NuMI-L-375, May 1998.




Figure 1: Correlations of atmospheric oscillation amplitudes: shown are contours of  $A_{atm}^{\mu e}$  in the  $(A_{atm}^{\mu\mu}, A_{atm}^{e\ell})$  plane for the (a) negative root, and (b) positive root solutions for  $P_{\mu 3}$  in Eq. (30), which correspond to  $\nu_\mu$  being more closely associated with  $\nu_2$  and  $\nu_3$ , respectively. Also shown are contours of  $A_{atm}^{e\tau}$  for the (c) negative root, and (b) positive root solutions. The point with bi-maximal solar and atmospheric neutrino oscillations is noted.